



پایان‌نامه‌ی کارشناسی ارشد: فریده ریگی، ۱۳۹۵

## بررسی پیرووات کیناز ایزوفرم M2 در بیماران مبتلا به سرطان کولورکتال و مقایسه آن با افراد طبیعی

سرطان کولورکتال، سومین عامل سرطان در سراسر جهان می باشد که بعد از سرطان ریه و پستان شایع ترین عامل مرگ و میر ناشی از سرطان می باشد. تومور مارکر M2-PK یک ایزوفرم آنزیم گلايکولیتیک PK می باشد که در سطح سلولهای در حال تکثیر سرطانی آشکار می شود و یک مارکر جدید برای تعیین سرطان کولورکتال می باشد. در این مطالعه نمونه مدفوع و پلاسمای 101 بیمار مراجعه کننده به بخش کولونوسکوپی قبل از انجام کولونوسکوپی گرفته شد که پس از انجام آزمایش M2-PK روی این نمونه هامشخص شد 68 نفر سالم، 12 نفر مبتلا به پولیپ و 21 نفر مبتلا به سرطان کولورکتال می باشند که این نتیجه با جواب پاتولوژی نمونه های ارسالی از بخش کولونوسکوپی به آزمایشگاه تائید شد. جهت اندازه گیری آنزیم M2-PK از کیت الیازی Biotech AG. SheBo استفاده شد که جهت اندازه گیری پلاسمای مدفوع دو کیت جداگانه طراحی شده است. در مقدار Cutoff برابر با 4 u/ml حساسیت تست M2-PK مدفوع در گروه نرمال و توموری 100% و ویژگی آن 66% محاسبه شد و در مقدار Cutoff برابر با 15 u/ml حساسیت تست M2-PK plasma در گروه نرمال و توموری 36% و ویژگی آن 55% برآورد گردید.

بنابراین با توجه به نتایج بدست آمده در این مطالعه که حساسیت و اختصاصیت بالای تست M2-PK مدفوع و پلاسمای بیماران مبتلا به سرطان کولورکتال نشان داد می توان از این تست به عنوان یک روش غیرتهاجمی و ابزار تشخیصی برای تعیین سرطان کولورکتال استفاده کرد.

**کلیدواژه‌ها:** سرطان کولورکتال، آنزیم M2-PK، مدفوع و پلاسمای Biotech AG. SheBo

شماره‌ی پایان‌نامه: ۱۲۷۳۰۵۶۰۹۴۱۰۲۵

تاریخ دفاع: ۱۳۹۵/۰۵/۱۸

رشته‌ی تحصیلی: زیست‌فناوری (بیوتکنولوژی)

دانشکده: علوم پایه

استاد راهنما: دکتر علی‌اکبر جنت‌آبادی

استاد مشاور: حمید رضا هاشمیان

### **M.A. Thesis:**

Study of pyruvate kinase M2 isoform in patients suffering colorectal cancer and comparing it with the normal.



Colorectal cancer is the third cause for cancer in the world, and after breast and lung cancers is the most common reason for death caused by a cancer. M2-PK tumor marker is an isoform from PK Glycolytic enzyme that appears in proliferating cancer cell and is a new marker to determine colorectal cancer. In this study, stool and plasma sample of 101 patients was taken before referring to colonoscopy section. After conducting the M2-PK experiment, it became evident that 68 were healthy, 12 were suffering polyps, and 21 were suffering colorectal cancer that was reassured by the pathology results of samples sent from colonoscopy section. To measure M2-PK enzyme, Biotech AG.SheBo ELISA kit was used; two different kits were used to measure plasma and stool. With 4uml Cutoff, sensitivity of M2-PK test was within normal group and 100% tumor with 66% specificity was calculated. And with 15uml Cutoff, sensitivity of M2-PKPlasma test was within normal group and 36% tumor and 55% specificity was estimated.

Therefore, according to the obtained results in this study, high specificity and sensitivity of M2-PK test of stool and plasma for patients suffering colorectal cancer show that this test can be used as a non-invasive method and a diagnostic tool in determining the colorectal cancer.